

Certificate of Analysis

PCR Nucleotide Mix:

Part No.	Size
C114G	200µl
C114H	1,000µl

Description: PCR Nucleotide Mix is a premixed solution containing the sodium salts of dATP, dCTP, dGTP and dTTP, each at a concentration of 10mM in water; the total concentration of nucleotides, therefore, is 40mM (pH 7.5). This solution is ready to use and is optimized for standard polymerase chain reactions and specialty approaches including hot-start and reverse transcription PCR (RT-PCR).

Storage Conditions: See the Product Information Label for storage recommendations. Avoid exposure to frequent temperature changes. Mix well prior to use.

Usage Note: Concentration gradients may form in frozen products and should be dispersed upon thawing. Mix well prior to use.

Quality Control Assays

Activity Assays

DNase and RNase Assay: To test for nuclease activity, 50ng of [³H]DNA or 50ng of [³H]RNA is incubated with 5µl of PCR Nucleotide Mix for 2 hours at 37°C in 1X MULTI-CORE™ Buffer (Cat.# R9991). The release of radiolabeled nucleotides is measured by liquid scintillation counting of TCA-soluble material. The specification is <3% release.

Physical Purity: The individual dNTPs contained in the PCR Nucleotide Mix each contain ≥99.0% triphosphate as measured by HPLC.

PCR Functional Assay: The individual dNTPs contained in the PCR Nucleotide Mix are tested using PCR to amplify a 360bp region of the α-1-antitrypsin gene. The amplification product is detected as a single 360bp band following agarose gel electrophoresis and ethidium bromide staining.

RT-PCR Functional Assay: The PCR Nucleotide mix is tested using RT_PCR to produce a 323bp amplicon from 0.25 zeptomoles of starting RNA template (100 copies).

PCR Nucleotide Mix

REF C1141 LOT 0000212857
-30°C to -10°C 2019-04-25
200µl Dispensed Lot#: 0000202721

For Laboratory Use

Country of Origin: USA

Promega Corporation
2800 Woods Hollow Road
Madison, WI 53711-5399 USA



ADC1141 00002128572

PEEL HERE



Promega

Promega Corporation

2800 Woods Hollow Road	
Madison, WI 53711-5399	USA
Telephone	608-274-4330
Toll Free	800-356-9526
Fax	608-277-2516
Internet	www.promega.com

PRODUCT USE LIMITATIONS, WARRANTY DISCLAIMER

Promega manufactures products for a number of intended uses. Please refer to the product label for the intended use statements for specific products. Promega products contain chemicals which may be harmful if misused. Due care should be exercised with all Promega products to prevent direct human contact.

Each Promega product is shipped with documentation stating specifications and other technical information. Promega products are warranted to meet or exceed the stated specifications. Promega's sole obligation and the customer's sole remedy is limited to replacement of products free of charge in the event products fail to perform as warranted. Promega makes no other warranty of any kind whatsoever, and SPECIFICALLY DISCLAIMS AND EXCLUDES ALL OTHER WARRANTIES OF ANY KIND OR NATURE WHATSOEVER, DIRECTLY OR INDIRECTLY, EXPRESS OR IMPLIED, INCLUDING, WITHOUT LIMITATION, AS TO THE SUITABILITY, PRODUCTIVITY, DURABILITY, FITNESS FOR A PARTICULAR PURPOSE OR USE, MERCHANTABILITY, CONDITION, OR ANY OTHER MATTER WITH RESPECT TO PROMEGA PRODUCTS. In no event shall Promega be liable for claims for any other damages, whether direct, incidental, foreseeable, consequential, or special (including but not limited to loss of use, revenue or profit), whether based upon warranty, contract, tort (including negligence) or strict liability arising in connection with the sale of the failure of Promega products to perform in accordance with the stated specifications.



PCR Satisfaction Guarantee

Promega's PCR Systems, enzymes and reagents are proven in PCR to ensure reliable, high performance results. Your success is important to us. Our products are backed by a worldwide team of Technical Support scientists. Please contact them for applications or technical assistance. If you are not completely satisfied with any Promega PCR product we will send a replacement or refund your account.
That's Our PCR Guarantee!

Product must be within expiration date and have been stored and used in accordance with product literature. See Promega Product Insert for specific tests performed.

Signed by:

J. Stevens

J. Stevens, Quality Assurance

© 1997–2011 Promega Corporation. All Rights Reserved.

MULTI-CORE is a trademark of Promega Corporation.

Triton is a registered trademark of Union Carbide Chemicals & Plastics Technology Corporation.

All specifications are subject to change without prior notice.

Product claims are subject to change. Please contact Promega Technical Services or access the Promega online catalog for the most up-to-date information on Promega products.

Part# 9PIC114
Printed in USA. Revised 10/11

1. Description

Applications of PCR Nucleotide Mix that have been tested include:

- PCR amplification by standard approaches.
- PCR amplification by specialty approaches such as hot-start and RT-PCR.

PCR Nucleotide Mix may be applicable to other specialty PCR approaches such as high-fidelity and long PCR.

2. Standard Applications

Preparing the Reaction Mix for PCR Using *Taq* DNA Polymerase

Reagents to Be Supplied by the User

(Solution compositions are provided in Section IV.)

- MgCl₂, 25mM (Cat.# A3511, A3512 or A3513)
- Nuclease-Free Water (Cat.# P1193)
- *Taq* DNA polymerase, 5u/μl
- 10X reaction buffer with 15mM MgCl₂
- 10X reaction buffer without MgCl₂ (optional)
- upstream and downstream primers, each at 20μM

1. Completely thaw, vortex and centrifuge all reagents before beginning the procedure.

Note: Optimal conditions, including reaction times, temperatures and reagent concentrations are dependent on the *Taq* DNA polymerase, template and primers used. The **magnesium concentration is especially important** and should be titrated between the range of 1.5–3.0mM in order to ensure optimal results. In many cases, a magnesium concentration of 1.5mM will result in satisfactory amplification. Therefore, two reaction preparations are given below. The first reaction is for 10X reaction buffer with 15mM MgCl₂. The second reaction is for 10X reaction buffer **without** 15mM MgCl₂.

2. Prepare one of the reaction mixtures listed below by adding reagents to a sterile microcentrifuge tube in the specified order.
 - a. Reaction components for **10X reaction buffer with 15mM MgCl₂**:

Component	Component Volumes	Final Concentration
10X reaction buffer (with 15mM MgCl ₂)	5μl	1X
PCR Nucleotide Mix (10mM each dNTP)	1μl	*800μM
upstream primer, 20μM	0.25–2.5μl	0.1–1μM
downstream primer, 20μM	0.25–2.5μl	0.1–1μM
<i>Taq</i> DNA polymerase, 5u/μl	0.25μl	0.025u/μl
template DNA	Xμl	<250ng
Nuclease-Free Water to final volume of	50μl	

- b. Reaction components for **10X reaction buffer without 15mM MgCl₂**:

Component	Component Volumes	Final Concentration
MgCl ₂ , 25mM	3μl	1.5mM
10X reaction buffer (without 15mM MgCl ₂)	5μl	1X
PCR Nucleotide Mix (10mM each dNTP)	1μl	*800μM
upstream primer, 20μM	0.25–2.5μl	0.1–1μM
downstream primer, 20μM	0.25–2.5μl	0.1–1μM
<i>Taq</i> DNA polymerase, 5u/μl	0.25μl	0.025u/μl
template DNA	Xμl	<250ng
Nuclease-Free Water to final volume of	50μl	

*The final concentration of each dNTP is 200μM.

3. Vortex the mixture and centrifuge briefly to ensure all liquid is at the bottom of the tube.
4. If using a thermal cycler without a heated lid, overlay the surface of the reaction mixture with 25–50μl of mineral oil.
5. Place the reaction tubes in a thermal cycler. The times and temperatures for denaturation, annealing and extension phases of amplification are template- and primer-dependent. Follow the general amplification guidelines in Section III as a starting point.

3. General Guidelines for Amplification by PCR

The following guidelines apply to target sequences between 200 and 2,000bp and are optimal for the Perkin-Elmer Thermal Cycler Model 480 or comparable thermal cyclers.

Note: Optimal denaturation and annealing reaction times for the Perkin-Elmer Thermal Cycler Model 9600, or comparable thermal cyclers, are shorter.

A. Denaturation

- Generally, a 2-minute initial denaturation step at 95°C is sufficient.
- Subsequent denaturation steps will be between 30 seconds and 1 minute.

B. Annealing

- Optimize the annealing conditions by performing the reaction starting approximately 5°C below the calculated melting temperature of the primers and increasing the temperature in increments of 1°C to the annealing temperature.
- The annealing step is typically 30 seconds to 1 minute.

C. Extension

- The extension reaction is typically performed at the optimal temperature for *Taq* DNA Polymerase, which is 72–74°C.
- Allow approximately 1 minute for every 1kb of DNA to be amplified.
- Minimum extension time should be 1 minute.
- A final extension of 5 minutes at 72–74°C is recommended.

D. Soaking

- If the thermal cycler has a "soak" cycle, the cycling reaction can be programmed to end by holding the tubes at 4°C for several hours.
- This cycle can minimize any polymerase activity that might occur at higher temperatures, although this is not usually a problem.

E. Cycle Number

- Generally, 25–30 cycles result in optimal amplification of desired products.
- Occasionally, up to 40 cycles may be performed, especially for detection of low-copy message.

4. Composition of Buffers and Solutions

10X reaction buffer with 15mM MgCl₂

100mM	Tris-HCl (pH 9.0 at 25°C)
500mM	KCl
15mM	MgCl ₂
1%	Triton® X-100

10X reaction buffer without 15mM MgCl₂

100mM	Tris-HCl (pH 9.0 at 25°C)
500mM	KCl
1%	Triton® X-100